

VIASYUK, P.A., otvetstvennyy red.; VASILENKO, A.A., red.; YUKHIMCHUK, F.F.,
kand.sel'skokhozyaystvennykh nauk, red.; ZELIGMAN, S.B., kand.
tekhn.nauk, red.; KUKHARENKO, N.I., kand.biol.nauk, red.;
MULYARSKIY, B.Ya., red.izd-va; SIVACHENKO, Ye.K., tekhn.red.

[Improving techniques of using fertilizers] Usovershenstvovanie
tekhniki vneseniia udobrenii. Kiyev, 1955. 255 p. (MIRA 11:6)

1. Akademiya nauk URSR. Kiyev. Rada po vyvcheniyu produktivnykh
sil URSR. 2. Deystvitel'nyy chlen Akademii nauk USSR i Vsesoyuznoy
akademii sel'skokhozyaystvennykh nauk im. V.I.Lenina (for Viasyuk)
3. Deystvitel'nyy chlen Akademii nauk USSR (for Vasilenko)
(Fertilizers and manures)

Vlasovuk P4

Action of nuclear radiations on plants. P. A. Vlasovuk
Sessiya Akad. Nauk S.S.R. po Mirnomu Teplofisicheskiyu MC
Akademii Energetiki 1955, Zasedaniya Otdel. Biol. Nauk, 127-47
(English summary, 147-8).—The use of Zn^{65} , Co^{60} , Ca^{45} ,
 S^{35} , P^{32} , and C^{14} at the av. activity of 0.01-100 μ c./kg. of
seeds of various plants resulted in the following observations
which supplement previous material which is reviewed.
Ionizing radiations cause a 2-phase response in plants:
the initial phase is that of suppression or of stimulation, the
2nd phase is that of normalization. Small doses of radio-
active materials introduced into seeds of planting soil led to
improved utilization of nutrients, accelerated metabolism,
and increased crops; as such this can be recommended for
agricultural use. Such mild treatment also acts as a form of
buffer against harmful effects of large doses of radiation;
this protective action is also shown by the SH compds. and
ascorbic acid in the plant tissues. The effect of ionizing
radiation on the plants is greatly affected by the nutritional
state of the plants. Expts. were run with yeast, wheat, flax,
and sugar beet, and numerous radiograms are shown.
G. M. Kostylev

Vlas'yuk, D. N.

No 4,

✓ The significance of different forms of phosphates in plant nutrition. P. A. Vlas'yuk, E. S. Kosmatyl, and Z. M. Klimovitskaya (Inst. Plant and Agrochem., Acad. Sci. Ukr. S.S.R., Kiev). *Fiziol. Rastenii, Akad. Nauk S.S.R.* 2, 334-7(1955).—Expts. with N-P-K plant diet contg. P^{32} in superphosphate, Ca pyrophosphate, or Ca orthophosphate were performed on sugar beet, wheat, and clover. Sugar beet and clover utilize the P content of superphosphate most intensely, pyrophosphate is utilized less well, and orthophosphate the least. P, regardless of its source, is localized more in the constitutional proteins than in protein reserves (storage proteins). In sugar-beet leaf in the 16-day plants (initial vegetative period) there are formed, in addition to inorg. P, glycerophosphate, glucose-1-phosphate, and fructose-1,6-diphosphate. At this age the roots show a considerable concn. of the inorg. P only. The best intake of P from superphosphate into sugar beet occurs when the fertilizer is introduced into the rows at planting. If the superphosphate is introduced some 5 cm. below the seeds the intake of P is considerably reduced. The meristematic young tissues of plants are richer in P than are the older tissues.

G. M. Korolapoff

(2)

VLASYUK P. A.

USSR.

[Participation of microorganisms in plant nutrition.] P. A. Vlas'yuk and V. D. Manzon (Inst. Plant Physiol. and Agro-chem., Acad. Sci. Ukr. S.S.R., Kiev). *Izvest. Akad. Nauk S.S.R., Ser. Biol.* 1955, No. 3, 49-53.—The use of S^{35} and P^{32} tracing by means of labeled phosphate and sulfate introduced into *Azotobacter* cultures, in connection with nutrition of corn and barley plants, conducted so that the sole source of labeled atoms was their contents in the bacterial bodies, showed that both live and dead bacterial material directly participate in plant nutrition. In sterile conditions P enters corn plants more readily from dead bacteria than from live ones; the same applies to barley. G. M. K.

Vlas'yuk, P. A.

MD

✓ The inactivation of urease by compounds of manganese as a preventative for the loss of urea nitrogen. P. A. Vlas'yuk and A. V. Mauorik. *Dopovidi Akad. Nauk Ukr. R.S.R.* No. 4, 364-7 (1955) (Russian summary).—Test materials were MnSO₄ and waste products of Mn mining. MnSO₄ was added at the rate of 0.25 (I) and 2.5 g./l. (II), and the waste material at the rate of 7.5 g./l. (III). Untreated urine was used as control (IV). Original urea-N content was 3.45 g.; 5 days after the addn. of the reagents urea N in IV was 1.20 and 10 days later, 0.60 g., or losses in N of 65.2 and 74.0%. In I the urea-N values were in 5 days 3.40 and in 10 days 3.38 g. or losses of 1.5 and 2.1%. In II urea N was in 5 days 3.39 and in 10 days 3.39 g. or a loss of 1.8 and 1.8%. In III urea N was in 5 days 3.42 and in 10 days 3.40 g. or a loss of 0.9 and 1.5%.

B. S. Levine

Inst. Plant Physiology & Agrochem, AS Ukr SSR

VLASYUK, P.A., MANRIK, A.V.

Increasing the biological activeness of the soil subjected to
enriched composts. Dop. AN URSR no.5: 500-504 '55. (MLRA 9:3)

1. Diysniy chlen AN URSR (for Vlasyuk); 2. Institut fiziologii
roslin ta agrokhimii AN URSR.
(Compost)

VLASYUK, P.A.

✓ 13544* Effect of Organo-Mineral Fertilizers on Increasing
the Yield of Corn. Vlijanie organo-minerálnykh udobrenii
na povyshenie urozhaja kukuruzy. (Russian) P. A. Vlasjuk AG
and P. Z. Liseval. Zemledelie, v. 3, no. 7, July 1953, p. 47-51.
Effect of $N_2P_2K_2$, K_2SO_4 , KCl , $K-Mg$, and Mn with or
without compost. Tables.

"APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7

VLASYUK, P.A.

✓ 2566. Metabolism in closer plants studied in the radioactive sulphur

Metabolism of radioactive sulphur in the plants
of the genus *Solanum* was studied by the method of protein fractionation.

APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7"

"APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7

✓ 6657. Phosphorus metabolism in the sugar beet. P. A. Vlasuk
and F. S. Krasnolutskii. *Vestn. Akad. Sel'sk. Nauk. UkrSSR.*, 1956, No. 1, p. 15811 - In the present work attention is given to the question of the metabolism of phosphorus in the sugar beet. The results show that

APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7"

VLAS YUK, P.A.

VLASYUK, P.A.; PORUTS'KIY, G.V.

Phytohormones and the vitality of a plant organism. Bot. zhur.
[Ukr.] 12 no.2:77-89 '55. (MIRA 8:10)
(Hormones (Plants))

VLASYUK, P.A.; LISOVAL, P.Z.; DOBROTVORS'KA, O.M.

Properties of organic and mineral composts and their effect on
the yield of sugar beets. Mikrobiol. zhur. 17 no.4:15-21 '55
(MLRA 10:5)

1. Z Institutu fiziologii roslin ta agrokhimii AN URSR
(COMPOST) (SUGAR BEETS)

"APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7

APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7"

"APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7

APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7"

VLASYUK, P.A.

✓ The influence of root and foliar feeding of radioactive phosphorus and calcium on the growth and sugar content of sugar beets. P. A. Vlasyuk, E. S. Kosmatyi, and Z. M. Klinovitskaya. Doklady Nauk. Akad. Sel'skokhoz. Nauk im. V. I. Lenina 10, No. 6, 15-17(1955).—Supplying to 16 kg. soil 5-10 microcuries of radionuclide Ca has increased the wt. of the roots and leaves by 21% and the sugar content by 0.3-0.8%. Radioactive phosphorus (in the form of Na_2HPO_4) applied to the foliage has increased the wt. of roots by 41%. All through the growing period there was more Ca in the leaves than in the roots. J. S. Joffe

VLASYUK, P.A.

813* Radioactive Isotopes and the Development of Plants.
Radioaktivnye izotopy i razvitiye rastenii. (Russian.) P. A. Vlasuk.
Nauka i zhizn, v. 22, no. 10, Oct. 1955, p. 21-22.
Effect of treating lupine seeds with ionizing radiations of Zn,
and of radioactive P isotopes on wheat seeds, etc.

VLASYUK, P.A.

Atomic energy in the service of agriculture. Visnyk AN URSR 26
no.9:35-41 S'55. (MIRA 8:11)

1. Diysniy chlen Akademii nauk URSR
(Radiobiology) (Agriculture)

VLASYUK, P.A.

I.V.Michurin's life and work; on the 100th anniversary of his birth.
Visnyk AN URSR 26 no.10:3-11 O '55. (MIRA 9:1)

I.Diysn.chlen Akademii nauk URSR i Vsesoyuznoy Akademii Sil'skogospo-
darchykh nauk imeni Lenina.
(Michurin, Ivan Vladimirovich, 1855-1935)

VLASYUK, P.A.

USSR/Biology - Plant physiology

Card 1/1 Pub. 22 - 51/53

Authors : Vlasyuk, P. A., Act. Memb. Acad. of Sc., Ukr. SSR.; and Grodzinskiy, D. M.

Title : Repeated utilization of phosphorus and sulfur by buckwheat

Periodical : Dok. AN SSSR 102/4, 845-847, Jun 1, 1955

Abstract : Biological data are presented on the re-utilization of P and S by buckwheat. Five references: 3 USSR and 2 USA (1932-1950). Tables.

Institution : Acad. of Sc., Ukr. SSR, Inst. of Plant Physiology and Agricultural Chem.

Submitted : January 8, 1955

VLASYUK, P.A.; GRODZINSKIY, D.M.

Tropisms of plant roots toward nuclear radiations. Dokl. AN SSSR 105
no.6:1358-1360 D '55. (MIRA 9:4)

1.Deystvitel'nyy chlen AN USSR (for Vlasyuk).2.Institut fiziologii
rasteniy i agrokhimii Akademii nauk USSR.
(Roots (Botany)) (Plants, Effect of radiation on)

~~Vlas'yuk~~ VLASYUK P.A.

96. Book Published on Uses of Radioisotopes in Studies of Plant Nutrition

Mikroelementy i Radioaktivnye Izotopy v Pitanii Rasteniy (Micro-elements and Radioactive Isotopes in Plant Nutrition), by Academician Petr Antipovich Vlasyuk, Academy of Sciences Ukrainian SSR, Kiev, Publishing House of Academy of Sciences Ukrainian SSR, 1956, 116 pp

Recent advances in scientific research in the fields of study and use of the microelements and radioactive substances in agriculture are discussed.

The chapter headings and pagination are as follows: foreword (3-4); improving conditions of plant nutrition by manganese fertilizers (5-21); superphosphate containing manganese -- a new type of fertilizer (22-26); significance of manganese microelement in increasing plant viability (27-38); selective biological properties of plants in relation to ultramicroelements (39-51); content of mobile forms of zinc, boron, cobalt, and copper micro-elements in soils of the Ukrainian SSR (52-59); use of tracer atoms for establishing methods for improving plant nutrition (60-75); effect of nuclear radiation on plants (76-90); effect of small doses of ionizing radiation from radioactive isotopes of zinc and cobalt on plants (91-104); and conclusion (105-114). (U)

ZINOV'YEVA, Khristina Gavrilovna; VLASYUK, P.A.; akademik, red.;
IGNATENKO, A.I., red.; KVITKA, S.P., tekhn. red.

[Azotobacter and farm plants] Azotobakter i sel'skokhoziaistven-
nye rasteniia. Kiev, Gos.izd-vo sel'khoz.lit-ry, USSR, 1962.
178 p.

(Azotobacter) (Field crops) (MIRA 16:3)

"APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7

VLASYUK, Petr Antipovich

[Principles of agriculture] Osnovy sil'skogo hospodarstva. Kyiv,
Radians'ka shkola, 1956. 532 p. (MIRA 10:11)
(Ukraine--Agriculture)

APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7"

VLASYUK, P.A.

[Ways of using radioactive isotopes in agriculture] Shliakhy
zastosuvannia radioaktivnykh izotopiv u sil's'komu hospodarstvi.
Kyiv, 1956. 59 p.
(Radioisotopes) (Agriculture)

(MLRA 10:4)

ALBRECHT, T. A.

SPIVAK, M.S., glavnnyy redaktor; BELOZUB, V.G., redaktor; VASILENKO, P.M., redaktor; ZORIN, I.G., redaktor; IL'CHENKO, I.K., redaktor; KOVAL', A.G., redaktor; KHYLOV, A.F., redaktor; PUHAL'SKIY, A.V., redaktor; SIDORENKO, A.P., redaktor; FEDCHENKO, A.N., redaktor; ANGELINA, P.N., redaktor; BUZANOV, I.F., redaktor; BOYKO, D.V., redaktor; BURKATSKAYA, G.Ye., redaktor; VASILENKO, A.A., redaktor; VIASYUK, P.A., redaktor; GORODNIY, N.G., redaktor; DEMIDENKO, T.T., redaktor; DUBKOVETS'KIY, F.I., redaktor; KIRICHENKO, F.G., redaktor; LITOVGHENKO, G.P., redaktor; OZERNYY, M.Ye., redaktor; PERSHIN, P.N., redaktor; POPOV, F.A., redaktor; POSMITNYY, M.A., redaktor; PSHEMICHNYY, P.D., redaktor; RADCHENKO, B.P., redaktor; ROMANENKO, I.N., redaktor; RUBIN, S.S., redaktor; SAVCHENKO, M.Kh., redaktor; SOKOLOVSKIY, A.N., redaktor; TSYBUNKO, K.Ye., redaktor; KOVAL'SKIY, V.F., tekhnicheskiy redaktor

[Practical collective farm encyclopedia] Kolkhoznaia proizvodstvennaia entsiklopediya. Izd.2-oe, ispr. i dop. Kiev, Gos.izd-vo sel'khoz. lit-ry USSR. Vol.1. Abrikos - liutserna. 1956. 688 p. (MIRA 10:9)
(Agriculture--Dictionaries)

SPIVAK, M.S., golovnyy redaktor; BILOZUB, V.G., redaktor; VASILENKO, P.M., redaktor; ZORIN, I.G., redaktor; IL'CHENKO, I.K., redaktor; KOVAL', O.G., redaktor; KRILOV, O.F., redaktor; PUHAL'S'KIY, A.V., redaktor; SIDORENKO, O.P., redaktor; FEDCHENKO, O.N., redaktor; ANGELINA, P.M., redaktor; BUZANOV, I.F., redaktor; BOYKO, D.V., redaktor; BURKATS'KA, G.E., redaktor; VASILENKO, A.O., redaktor; VIASYUK, P.A., redaktor; GORODNIY, M.G., redaktor; DEMIDENKO, T.T., redaktor; DUBKOVETS'KIY, F.I., redaktor; KIRICHENKO, F.G., redaktor; LITOVCHENKO, G.P., redaktor; OZERNIY, M.O., redaktor; PERSHIN, P.M., redaktor; POPOV, F.A., redaktor; POSMITNIY, M.O., redaktor; PSHENICHNIY, P.D., redaktor; RADCHENKO, B.P., redaktor; POMANENKO, S.S., redaktor; RUBIN, S.S., redaktor; SAVCHENKO, M.Kh., redaktor; SOKOLOVS'KIY, O.N., redaktor; TSIBENKO, K.O., redaktor; SHCHERBINA, O.P., redaktor; KRAVCHENKO, M.F., tekhnichniy redaktor

[Collective farm encyclopedia] Kolhospna vyrobnycha ensyklopediia.
Vyd. 2-e, perer. i dop. Kyiv, Derzh.vyd-vo sil's'kohospodars'koi
lit-ry URSS. Vol.1. Abrykos - Liutserna. 1956. 756 p. (MIRA 9:9)
(Agriculture--Encyclopedias and dictionaries)

USSR/Cultivated Plants. Cereals.

M

Abs Jour: Ref Zhur-Biol., No 17, 1958, 77625.

Author : Vlas'yuk P. A.; Lisoval, P.Z.

Inst : AS UkrSSR.

Title : Influence of Organic-Mineral Fertilizers on
Harvests of Corn.

Orig Pub: V sb.: Vopr. razvitiya s.-kh. Nauk. Kiev,
AN USSR, 1956 (1957), 40-49.

Abstract: In the Institute of Physiology of Plants and
Agrotechny AS UkrSSR in 1953-1954, the influ-
ence was studied of different types of ferti-
lizers on harvests of corn, placed in crop-rota-
tion after winter wheat. On meadow-chernozem
soils, organic-mineral fertilizers contributed
to the increase of the grain harvest of corn by

Card : 1/3

USSR/Cultivated Plants. Cereals.

M

Abs Jour: Ref Zhur-Biol., No 17, 1958, 77625.

14.1-15.7 c/ha with harvests in the control (without fertilizer) 39.1 c/ha. Application of 5 t/ha of organic-mineral composts in a nest with sowing proved the most effective and assured on weakly-podzolic, sandy soils the obtaining of a grain harvest of corn of 81.6 c/ha with harvest in the control 39 c/ha. On weakly podzolic, lightly sandy soils the application of 5 t/ha of humus in the nest with sowing assured the addition of harvest of 7 c/ha with harvest in control 42.1 c/ha. With an organic-mineral mixture with the same control/grain-harvest increase of 24 c/ha was obtained. Tillage of green mass of harvested lupine with combined

Card : 2/3

USSR/Cultivated Plants. Cereals.

M

Abs Jour: Ref Zhur-Biol., No 17, 1958, 77625.

application of 3 t/ha of organic-mineral mixture
in a nest on weakly podzolic sandy soil gave a
harvest of green mass of corn of the Odessa 10
variety in 1954 of 1087 c/ha. -- N. P. Fedorova.

Card : 3/3

38

USSR/Soil Science. Organic Fertilizers.

J-4

Abs Jour: Ref Zhur-Biol., No 6, 1958, 24759.

Author : Vlasyuk, P.A.; Manorik, A.V.

Inst :

Title : Utilization of Enriched Composts for Increasing
Harvests of Agricultural Crops.

Orig Pub: V sb.: Vopr. razvitiya s. kh. Poles'ya Kiyev, AN
USSR, 1956 (1957), 73-80.

Abstract: The Institute of Plant Physiology and Agricultural Chemistry AN USSR utilized wastes of the brown coal industry for making composts with manure; content 0.53% N, 0.63% P, and 0.39 K₂O. In experiments for 6 mos. of 1951-1952 with storing of manure without the wastes of brown coal industry, the average

Card : 1/2

23

USSR/Soil Science. Organic Fertilizers.

J-4

Abs Jour; Ref Zhur-Biol., No 6, 1968, 24759.

losses of N constituted 13%, while, when compost was made with the wastes, the average content of N increased by 16.5%. The content of hydrolyzed N and of fixed ammonium was considerably increased by activity of compost. In the 1954 experiments on the application of P³², it was found that phosphorus from phosphorites turns to forms utilizable by plants. The increase of the sugar beet harvest due to the making a compost of manure with mineral fertilizers and brown coal constituted 69 c/ha., with a yield of 339 c/ha without addition of brown coal in the compost. Making a compost of manure with wastes of the brown coal industry jointly with phosphorus fertilizers and wastes of the manganese-ore industry proved most effective.

Card : 2/2

P
VLASYUK, V.A.; MANORIK, A.V.

Increasing peat efficiency by strengthening its biological activity.
Dop. AN URSR no. 1:79-84 '56. (MIRA 9:7)

1. Diysniy chlen AN URSR i VASGNIL (for Vlasyuk). 2. Institut fiziolegii
reslin ta agrokhimii AN URSR.
(Peat)

VLASYUK, P.A.

USSR/Soil Science - Physical and Chemical Properties of Soils. J-2

Abs Jour : Ref Zhur - Biol., No 2, 1958, 5767

Author : Vlasyuk, P.A.

Inst : Academy of Sciences LatvSSR

Title : The Content of Extractable Forms of Zinc, Boron, Cobalt, and Copper in the Soils of the Ukrainian SSR.

Orig Pub : Mikroelementy v s. kh. i meditsine, Riga, Akad Nauk LatvSSR, 1956, 97-103

Abstract : The microelement content of the turf-podzolic soils, gray forest soils, chernozems, solonetz and solonchak soils of Poles'ye, the right- and left-bank soils of the (Losostep') Wooded Steppe, the southern steppe regions and the Trans-Carpathian region of the UkrSSR is examined. The content of extractable zinc in the soils reflects its content in the soil-forming plants, while its distribution in the soil

Card 1/3

USSR/Soil Science - Physical and Chemical Properties of Soils.

J-2

Abs Jour : Ref Zhur - Biol., No 2, 1958, 5767

is found mainly in the horizons below the tillable level, while with the chernozems it is found in the tillable horizon. In the turf-podzolic and gray forest soils there is 0.23-0.43 mg. of Co per kilogram of soil (dry weight). In some gray forest soils and podzolized chernozems the Co content reaches 0.6 mg., and in the meadow-chernozem cultivated soils it reaches 2.3 mg. Most of the soil variations in the UkrSSR, with the exception of the peat and turf-podzolic soils, are fully supplied with Cu. The turf-carbonate and mountain meadow soils of Trans-Carpathia, and also the solonetz soils, contain especially large quantities of it; the chernozems and dark chestnut soils contain less, while the gray forest soils and turf-podzolic soils contain the lowest amounts. The Cu is most extractable in all types of soil where the mechanical composition is light, less so in heavy argillaceous soils, and least of all in peat soils.

Card 3/3

USSR / Soil Science. Mineral Fertilizers.

J-4

Abs Jour: Ref Zhur-Biol., No 8, 1958, 34425.

Author : Vlasvuk, P. A.

Inst : AS LatvSSR - Institute of Physiology of Plants
and Agrochemistry of AS UkrSSR.

Title : Improvement of Conditions of Nutrition of Plants
by Ions of Manganese Trace Fertilizers.

Orig Pub: V sb.: mikroelementy v s.kh. i meditsine, Riga,
AN LatvSSR, 1956, 111-124.

Abstract: According to numerous laboratory tests and field experiments, the positive action of Mn on respiration, photosynthesis, activity of enzymes, content of chlorophyll, etc., has been shown; in this connection, a considerable increase in yield and quality has been obtained in the following plants: sugar beets, winter wheat, corn, oats,

Card 1/2

43

USSR / Soil Science. Mineral Fertilizers.,

J-4

Abs Jour: Ref Zhur-Biol., No 8, 1958, 34426.

Abstract: buckwheat, millet, tobacco, potatoes, cucumbers, tomatoes, cabbage, eggplant, hemp, flax, garden strawberry and strawberry. Increase of the action of nitrification and ammonification by bacteria in the soil, as well as the inactivation of the ferment of urease in the liquid manure, has been established; in this connection, the decomposition of the urea is removed, and losses of N are sharply lowered. The yield of agricultural plants is particularly well affected by manganeseized P_s and by manganese slag. The task has been accomplished in the Institute of Physiology of Plants and Agrochemistry of the Academy of Sciences UkrSSR.
-- A. P. Shchorbakov.

Card 2/2

"APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7

APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7"

VLASYUK, P. H.

A S Ukr 558

Polar differentiation and alteration of stage readiness of plants for flowering. P. A. Vlasuk and G. V. Porutskii (Plant Physiol. Agrobiol. Inst., Kiev). *Izvest. Akad. Nauk S.S.R., Ser. Biol.* 1956, No. 1, 84-90.—The studies with cotton plants (high-Mn requirement) and corn (low-Mn requirement) were made in respect to content of dry matter and H₂O content in the progressive segments of the growing plants at various stages. Vernalization temp. limits reach 0° for cotton and 40° for corn. Vernalizing activity of MnSO₄ is more pronounced in cotton than in corn. Mn supply tends to increase the height periodicity of the content of dry matter and polarity in cotton, while the vernalization temp. is more important for corn. The polar periodicity of distribution of matter in the stems can be used as an index of the plant's readiness for flowering. G. M. Kosolapoff

MD

(1)

V. L. Vlas'yuk, Ph.D.

USSR/General Division - General Problems. Philosophy.
Methodology.

A-1

Abs Jour : Ref Zhur - Biologiya, No 7, 10 April 1957, 25639
Author : Vlas'yuk, P.A.
Inst : Academy of Sciences UkrSSR: Inst. of Agric. Biology; Inst.
of Plant Physiology and Agric. Chemistry; Inst. of
Entomology and Phytopathology.
Title : Increasing the Role of Agricultural Science in the
Development of Agriculture in the Ukraine.
Orig Pub : Visnik AN URSR, 1956, No 2, 27-38

Abst : The Academy of Sciences UkrSSR is studying a number of im-
portant problems connected with increasing crop yields
and improving productivity in animal farming in the light
of the directives of the XXth Congress of the CP USSR.
The Institute of Plant Physiology and Agricultural Chemis-
try has developed a number of new fertilizing techniques
applicable to various areas of the Ukraine and variety of

Card 1/2

USSR/General Division - General Problems. Philosophy.
Methodology.

A-1

Abs Jour : Ref Zhur - Biologiya, No 7, 10 April 1957, 25639

crops. 1955 marked the beginning of studies in the USSR of a new type of fertilizer: manganated granular super-phosphate, which increases potato yield by 24 cent/hect, sugar beet yield by 20 cent/hect, corn yield by 6 cent/hect, etc. The Institute of Entomology and Phytopathology is developing and putting into use new techniques in fighting pests and weeds. The Institute of Agricultural Biology is engaged in developing new crops. Soil scientists are working on problems of agricultural technique and fertilizers. Extensive use is being made of tracer element techniques. Ukrainian scientists have assumed the responsibility of pursuing further the theoretical problems connected with the use of atomic energy in farming.

Card 2/2

Vlas'yuk /A.
USSR/Microbiology. Soil Microbiology

F-3

Abs Jour : Ref Zhur-Biologiya, No 1, 1957, 576

Author : P. A. Vlas'yuk and V. D. Manzov
Inst :

Title : On the Application of Azotobacter for the
Enrichment of Composts with Atmospheric
Nitrogen

Orig Pub : Agrobiologiya, 1956, No 2, 89-97

Abstract : It was established that azotobacter when
introduced into composts (with phos-
phorite flour) of manure and straw, and
into composts made of peat became well
adjusted, absorbed nitrogen from the
atmosphere, and enriched with it the
medium. In the presence of azotobacter
losses of nitrogen from the composts of

Card 1/2

USSR/Microbiology. Soil Microbiology

F-3

Abs Jour : Ref Zhur-Biologiya, No 1, 1957, 576

Abstract : manure dropped from 37.8 to 18.9%,
and in composts from straw from 28.2 to
11.3%. In peat composts the nitrogen
content as compared with the original
increased from 3.2 to 10.7%, depending
on the composition of the composts. On
the introduction of the composts in-
filtrated with azotobacter into the soil,
the number of azotobacter in the soil
increased and displayed great activity.

Card 2/2

"APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7

APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7"

USSR/Cultivated Plants - Grains.

M-4

Abs Jour : Ref Zhur - Biol., No 9, 1958, 39235

Author : Vlas'yuk, P.A., Poruts'kiy, G.V.

Inst : AS UkrainianSSR.

Title : The Importance of Manganese Nutrient for Corn Growing
in the Ivales'ye Rayons of the UkrainianSSR.

Orig Pub : Nauk. pratsi vid. sil's'kozosp. nauk. AN URSR, 1956, vyp.
4, 4-15.

Abstract : Pre-sowing treatment of corn seeds with warm solutions of
manganese salts considerably increases the vitality, yield and
productivity of the plants. It contributes to the strengthen-
ing of the plant's metabolism and accelerates the
passage through the initial stages of growth and develop-
ment of the organism of the plant. -- Yu.P. Savchenko.

Card 1/1

- 46 -

VLASYUK, P.A., akademik.

Effectiveness of organomineral mixtures. Zemledelie 4 no.8:65-72
Ag '56.
(MLRA 10:1)

1. Institut fiziologii rasteniy Ukrainskoy SSR.
(Ukraine--Fertilizers and manures)

USSR/Cultivated Plants - Potatoes. Vegetables. Melons.

M-3

Abs Jour : Ref Zhur - Biol., No 7, 1958, 29794

Author : Vlasuk, P.A., Dolya, V.S.

Inst : Institute for Plant Physiology and Agrochemistry of the Academy of Sciences, Ukrainian SSR.

Title : The Effect of Micronutrients and Bacterial Fertilizers on the Output of Vegetable Pot Cultures.

Orig Pub : Dopovidi AN UkrSSR, 1956, No 6, 584-587 (ukr.; rez. russk.)

Abstract : It has been established as a result of experiments made by the Institute for Plant Physiology and Agrochemistry of the Academy of Sciences Ukrainian SSR in 1954-1955 that the application of micronutrients and phosphorus bacteria during the sprouting period considerably increases the growth of the vegetable cultures, shortens the budding time, that of flowering, of fruit ripening and

Card 1/2

USSR/Cultivated Plants - Potatoes. Vegetables. Melons.

Abs Jour : Ref Zhur - Biol., No 7, 1958, 29794

M-3

of cabbage head setting. The largest increase to the tomato yield was observed when phosphorus bacteria and manganese micronutrients were applied. Mn (the yield increased by 59 centners per ha.), Co (by 32 centners per ha.) and Zn (by 21 centners per ha.) proved most effective when raising cabbage sprouts in peat-compost pots.

Card 2/2

- 11 -

USSR/Soil Science. Mineral Fertilizers.

J-3

Abs Jour: Ref Zhur-Biol., No 6, 1958, 24746.

Author : Vlasyuk, P.A.; Lisovaya, P.Z.

Inst :

Title : Application of Kaluszite for Clover and Sugar Beet.

Orig Pub: Udobreniye i urozhay, 1956, No 11, 46-49.

Abstract: In the Institute of Physiology of Plants and Agricultural Chemistry of the AN USSR, in vegetating experiments on meadow-chernozem podsolic soil with red clover in its second year, the best results were given by: potassium sulfate, a mixture of it with potassium chloride and kaluszite. Increases of the yields comprised respectively 110; 88 and 77% in comparison with the baseline contained 16.0% K₂O; 5.5% MgO and 5.0% Cl. In the experiments with the

Card : 1/2

20

USSR/Soil Science. Mineral Fertilizers,

Abs Jour: Ref Zhur-Biol, No 6, 1958, 24746,

J-3

sugar beet, potash magnesia gave an increase of
16%, while kaluszite - 144%.

Card : 2/2

VLASYUK, P.A.

UKRAINE/Cultivated Plants - General Problems.

L-1

Abs Jour : Ref Zhur - Biologiya, No 16, 25 Aug 1957, 69179
Author : Vlasuk, P.A.
Inst :
Title : Basic Methods of Increasing Soil Fertility.
Orig Pub : Visnik AN URSR, 1956, No 12, 16-24

Abstract : Some measures of increasing soil fertility in the steppe
forestry zone of Ukrainian SSR are indicated (introduction
of fieldgrass rotation and use of fertilizers).

Card 1/1

USSR/Plant Physiology - General Problems.

Abs Jour : Ref Zhur .. Biol., No 18, 1958, 81966

I.

to the different duration of development stages (vernali-
zation, luminous). The advent of stage readiness in
the plants which were studied changed under the influence
of various conditions of growth (temperature, food regi-
men). This reflected itself in their polarity. Accord-
ing to the authors, the polar and layer differentiation
of the examined biochemical and physiological indexes can
serve as a criterion of the stage readiness of plants and
of the intensity of blossoming processes.
Bibliography, 43 titles. -- G.V. Porutskiy

Card 2/2

- 5 -

VLASTYUK, P.A.; PORUTSKIY, G.V.

On the results of the discussion on phytohormones. Ukr.bot.zhur.13
no.4:100-102 '56. (MIRA 10:1)
(Growth promoting hormones)

"APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7

VLASYUK, P.A.

Michurin's followers in the Ukraine. Nauk. zap. Kyiv. un. 15
no.11:31-34 '56. (MIRA 11:5)
(Ukraine--Agriculture)

APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7"

"APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7

VIA C P A

APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7"

"APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7

APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001860320004-7"

Effects of organic-inorganic fertilizers on microorganisms processes in the soil. P. A. Vlasov, K. M. Dobrotworskaya, and R. N. Olsuf'ev (State Univ., Kiev). Mikrobiologiya 25, 293-8 (1966).—Manure at 20 tons/ha. activated cellulose aerobes and N-fixing and nitrifying anaerobes in a winter wheat land more strongly than did a peat-compost blend at 3 tons/ha., even when reinforced with superphosphate (1.5 cvt) and lime (3 cvt). The intensification of ammonification, nitrification, and urea formation was also greater, so was the rise in wheat yield. Julian E. Smith

3

VLASYUK, P.A.

8146

ON THE EFFECT OF NUCLEAR RADIATION OF PLANTS.
P. A. VLASYUK and D. M. GRODZINSKII (Inst. of Plant Physiology and Agrochemistry). Doklady Akad. Nauk S.S.R. 106, 582-4 (1956) Jan. 21, (in Russian)

Studies of radioactive isotope effects on plants revealed many complicated and varying reactions produced by ionizing radiation. In some cases the same dose showed contrasting effects or no effects at all, depending on the temperature, nutrition level, or other external conditions. As a rule, the positive effects were produced by the small doses. Ionizing radiation produced two-phase reactions; small radiation doses intensified the physiological functions of the plant; the large doses slowed down and weakened the life functions of the cells and tissues, which eventually became restored to normal activity. Such two-phase reactions were observed in the study of Ca^{45} and P^{32} radiation on Saccharomyces cerevisiae. A similar parabolic reaction was observed in Uromyces corynephidis. A diagram is presented showing the effects of ionizing radioactive isotope Ca^{45} on the breathing functions of Saccharomyces cerevisiae culture (in % to the breathing intensity in stable isotope media). Experiments were carried out on sugar beet leaves and oat seeds. On the basis of the studies it may be concluded that the effects of ionizing radiation are determined by the functional conditions of the growing organism or by some of its separate organs. It is also possible that the radiation affects plants as background excitation. (R.V.J.)

2
Mos

VLASYUK P. A.

✓ 8147

EFFECTS OF SMALL DOSES OF IONIZING RADIATIONS
ON OXIDATION-REDUCTION PROCESSES IN PLANTS. P.
A. Vlasuk, Z. M. Klimovitskaya, and E. S. Kosmatif
(Ukrainian Inst. of Plant Physiology and Agrochemistry).
Doklady Akad. Nauk S.S.R. 106, 731-4 (1956) Feb. 1. (In
Russian)

Tracer studies of radioactive Ca^{45} (5 to 10 μ per plant)
effect on sugar beet yield and the effects of small doses of
 pu , Ca^{45} , and S^{35} in the food supply of sugar beets and
clover are tabulated and discussed. The studies established
that in the early stages of growth of sugar beets the ionizing
radiation increased the oxidation and lowered the reduction
processes. Considerable increase of reduction processes
over the oxidation which resulted in richer sugar beet yield
were observed towards the end of the vegetative period.

Larger ionizing radiation doses in clover plants increased
oxidation and suppressed the reduction process. In clover
plants small doses of radioactive S^{35} have intensified the ox-
idation, the tissue Iodo-reducing properties, and the content
of reducing forms of ascorbic acid. (R.V.J.)

3

Rec'd

U.S. Army K. H.

SPIVAK, M.S., glavnny red.; BELOZUB, V.G., red.; VASILENKO, P.M., red.; ZORIN, I.G., red.; IL'CHENKO, I.K., red.; KOVAL', A.G., red.; KRYLOV, A.F., red.; PUKHAL'SKIY, A.V., red.; SIDORENKO, A.P., red.; FEDCHENKO, A.N., red.; ANGELINA, P.H., red.; BUZANOV, I.F., red.; BOYKO, D.V., red.; BURKATSKAYA, G.Ye., red.; VASILENKO, A.A., red.; VLASYUK, P.A., red.; GORODNIY, N.G., red.; DEMIDENKO, T.T., red.; DUBKOVETSkiy, F.I., red.; KIRICHENKO, F.G., red.; LITOVCHENKO, G.P., red.; OZERNYY, M.Ye., red.; PERSHIN, P.N., red.; POPOV, F.A., red.; POSMITNYY, M.A., red.; PSHENICHNYY, P.D., red.; RADCHENKO, B.P., red.; ROMANENKO, I.N., red.; RUBIN, S.S., red.; SAVCHENKO, M.Kh., red.; SOKOLOVSKIY, A.N., red.; TSYBENKO, K.Ye., red.; KOVAL'SKIY, V.F., tekhn.red.

[Practical collective farm encyclopedia] Kolkhoznaia proizvodstvennaia entsiklopediia. Izd. 2-oe, perer. i dop. Kiev, Gos. izd-vo sel'khoz. lit-ry USSR. Vol.2. Malina-Iashchur. 1957. 923 p.

(Agriculture--Dictionaries)

(MIRA 11:4)

USSR / Soil Science. Mineral Fertilizers.

J-4

Abs Jour: Ref Zhur-Biol., No 8, 1958, 34385.

Author : Vlas'yuk, P. A., Lisoval, P. Z.
Inst : AS UkrSSR.

Title : Increase of Fertility of Soil and Yield in Agricultural Cultivations by Means of Utilizing Organic and Mineral Fertilizers.

Orig Pub: V sb.: Mestn. organ. udobreniya USSR, Kiyev,
AN USSR, 1957, 5-17.

Abstract: Based on experiments conducted on turf-podzolic sandy loam and meadowy-black earth podzolized soils, the authors claim that application of the organic-mineral system of fertilization secures the most favorable condition for the nutrition of plants and considerably increases the yield. The highest increase in yield was observed by

Card 1/2

USSR / Soil Science. Mineral Fertilizers.

J-4

Abs Jour: Ref Zhur-Biol., No 8, 1958, 34385.

Abstract: introduction of manure, mixed with P₂O₅ and defecation, The effectiveness of action of organic-mineral mixtures increases by adding manganese and lime residues to it, and also by composting organic fertilizers with the mineral. --
L. N. Kudryasheva.

Card 2/2

25

COUNTRY : USSR
CATEGORY : PLANT PHYSIOLOGY. Mineral Nutrition.
ABS. JOUR. : REF ZHUR. BIOLOGIYA. NO. 4, 1954. No. 15292
AUTHOR : Vlasyuk, P.A.
INST. : --
TITLE : Spectral Analytic Study of Selective properties in plants.
ORIG. PUB. : V sb.: Primeneniye metodov spektroskopii v prom-sti prodovol'stv. tovarov i s. kh., L., LGU, 1957, 51-59
ABSTRACT : Results are reported on a study by the emission-spectral method of the content of macro-, micro-, and ultramicroelements (34 elements in all) in various soils and plants cultivated in different zones of the USSR. Vegetable cultures selectively absorbed and accumulated (especially in the seed) a notable amount of microelements, although they were not successfully detected in the soils. Thus, cucumbers always contained Ag, Mo, Ib,

CARD: 1/2

2

COUNTRY

CATEGORY PLANT PHYSIOLOGY.

ABS. JOUR. REF ZHUR - BIOLOGIYA, NO. 4, 1959.

AUTHOR

INST.

TITLE

No. 15292

ORIG. PUB. :

ABSTRACT : Sn, V, and Cu; garlic - V; peppers - Mo, tomatoes - Zr; cabbages - I and Co; corn kernels - Au. The nature of the soil influenced the number of micro- and ultramicroelements in plants. -- N.I. Borisova

CARD:

2/2

COUNTRY : USSR
CATEGORY : Soil Science. Organic Fertilizers.
ABR. JOUR. : Kharkov., №. 43 1958, №. 104483
AUTHOR : Vlas'yuk, P. A.
TYPE: Kharkov University
TITLE : Improving Plant Nutrient Conditions With Brown Coal Tailings
OPIN. PER. : V sb.: Guminovyye udobreniya. Khar'kov, Khar'kovsk. un-t, 1957, 127-144
ABSTRACT : In absorptive power, brown coal tailings surpass a number of investigated substances and are second only to peat; the greatest absorption of P was observed for coal tailings of Khustskoye and Yurkovekoye origin; of ammonium, for Yasenovskoye brown coal tailings. The use of brown coal tailings together with mineral fertilizers or manure increases the yield of sugar beet, winter wheat, oats, rye, flax, corn and other crops, and noticeably increases the number of nodules on pea. The composting of manure with brown coal tailings caused a 50% decrease in P loss and an increase in the content of an active form of N of up to 117 mg per 100 g of

Card: 1/2

COUNTRY :	
CATEGORY :	J
AD. OF PUBL. :	RZhNKhCh., No. 23, 1992, No. 1044CZ
AUTHOR :	
INST. :	
TITLE :	
ORIG. SUB. :	
ABSTRACT :	soil as opposed to 78.8 mg without tailings; best results were obtained when composting manure with 10% brown coal tailings; an increase of the dose to 20% did not increase the effect. The addition of tailings to P when introducing it into the pits increased the cob harvest ^s of corn from 112.5 to 146.5 centners/hectare; the addition of coal tailings to K for flax cultivation increased the fiber yield from 7.4 to 10.3 centners/hectare and, besides, raised the oil content in the seeds from 43.2 to 53.5%.--S. A. Mamizov

Card: 2/2

J-4

USSR / Soil Science. Mineral Fertilizers.

Abs Jour: Ref Zhur-Biol., No 8, 1958, 34409.

Author : Vlasyuk, P. A.

Inst : Not given.

Title : New Aspect of Fertilization - Manganese Superglucophosphate.

Orig Pub: Byul. po fiziol. rasteniy, 1957, No 1, 6-10.

Abstract: Manganese Ps is obtained by means of granulation of Ps in powder form with 10 - 15% of manganese slime and 1 - 4% chalk. The fertilizer contains 17.8 to 18.3% of assimilated phosphoric acid, 2-3% of manganese oxides, and its free acidity is 1.7 - 2.7 (instead of 5% in Ps). According to the results of the tests, carried out by exper-

Card 1/2

USSR / Soil Science. Mineral Fertilizers.

J-4

Abs Jour: Ref Zhur-Biol., No 8, 1958, 34409.

Abstract: mental stations, as well as production experiments in collective farms of USSR, the efficiency of manganeseized P₂O₅ exceeds that of the common granulated P₂O₅. -- T. L. Rivkind.

Card 2/2

33

USSR / Plant Physiology. Mineral Nutrition.

I-3

Abs Jour: Ref Zhur-Biol., 1958, No 16, 7257'.

Author : Vlasyuk, P. A.; Manorik, A. V.
Inst : Ukrainian Scientific-Research Institute of Plant
Physiology.
Title : Admission of Radioactive P^{32} , S^{35} and C^{14} in Plants
from the Organic and Mineral Forms of Their Com-
pounds.

Orig Pub: Byul. po fiziol. rasteniy, 1957, No 1, 20-23.

Abstract: Winter wheat plants were fed with marked P_C in the
phase of tubing; in 7 days they were unearthed and
kept for two days in containers with water con-
taining 2 mcuries of P^{32} in the form of $Na_2HP^{32}O_4$.
The parts of the plants above ground were used as
green fertilizer under buckwheat. Bogatyr' buck-

Card 1/3

USSR / Plant Physiology. Mineral Nutrition.

I-3

Abs Jour: Ref Zhur-Biol., 1958, No 16, 72577.

Abstract: wheat was raised in soil cultures: P³² was introduced in the form of P_c, pure vegetative mass or infected with fresh manure. The accumulation of the raw mass and the weight of seed harvest were greatest in the case of application of mineral P, the least - in those fertilized only with vegetative residue. The admission of P³² with fertilization by vegetative mass without infection was more intensive than with it. During the cultivation of buckwheat on Knop's nutritive mixture, the rate was studied of the admission of S³⁵ (in the form of vitamin B₁ and methionine), C¹⁴ (in the form of tyrosine and Na₂C¹⁴O₃) and P³² (Na₂HP³²O₄). In the blossoming phase, a rapid admission of organic compounds was observed. Ten minutes after the appli-

Card 2/3

11

USSR / Plant Physiology. Mineral Nutrition.

I-3

Abs Jour: Ref Zhur-Biol., 1958, No 16, 72577.

Abstract: cation of tyrosine in a nutritive solution, the radioactivity of the flowers was highest; in 20-60 minutes, the radioactivity of the leaves was increased, but that of the flowers decreased. In three hours after the application of radioactive substances on the leaves, significant activity of the roots was detected through which methionine moved more intensively than vitamin B₁, and the NO₂CO₃ moved more intensively than the tyrosine. The work was carried out in the Ukrainian Scientific-Research Institute of Plant Physiology. --
B. Ye. Kravtsova.

Card 3/3

COUNTRY	USSR	
CATEGORY	:	N
ABS. JOUR.	RESiol., No.12,	1958, No.53935
AUTHOR	:	
TYPE	:	
FILED	:	
ORIG. PUB.	:	
ABSTRACT	wheat and oat yields following this were not noted, and a certain yield boost was observed in the test with oats. --N.I. Borisova	
CARD:	2/2	
6		

VLASYUK, P.; ZAKHARCHUK, P.; KALYUZHNYY, V.; PERESYPKIN, V.

Seventieth birthday of Mikhail Mikhailovich Godlin. *Pochtovedenie*
no.3:117-118 Mr '57. (MLRA 10:7)
(Godlin, Mikhail Mikhailovich, 1886-)

I.

USSR/Plant Physiology - Mineral Nutrition.

Abs Jour : Ref Zhur - Biol., No 23, 1958, 104360

Author : Vlasyuk, P.A., Kosmatyy, Ye.S., and Klimovitskaya, G.I.

Inst : Institute of Plant Physiology and Agrochemistry, AS
Ukrainian SSR.

Title : The Effect of Nitrato-Ammoniacal, Nitrogenous and Mineral
nous Nutrition on Sulfur Metabolism in the Sugar Beet.

Orig Pub : Fiziol. Rasteniy, 4, No 5, 432-439, 1957.

Abstract : Under conditions of a soil culture and a NPK background,
with respect to the sugar beet and wheat, it was estab-
lished through introducing $\text{Na}_2\text{S}^{35}\text{O}_4$ (50 curies per 16 kg
of soil) that, in contrast with P, more S enters into re-
serve proteins than into the constitutional proteins.
Injection into the roots of the sugar beet of aqueous so-
lutions of methionine or vitamin b₁ containing S³⁵ caused

Card 1/3

- 8 -

USSR/Plant Physiology - Mineral Nutrition.

I.

Abs Jour : Ref zhur - Biol., No 23, 1950, 104360

an intensive translocation of S into leaves, especially the younger leaves, in which connection a major part of S was also included in the reserve proteins. The rate of S metabolism (as determined according to its specific activity and the number of the individual forms of S) from methionine was higher than from vitamin B₁, especially with respect to the easily detached S fraction (by Shul'ts' method). A larger quantity of S entered into organic compounds than into mineral compounds. Compared with ammoniacal nutrition, the nitrate nutrition of the sugar beet favored an increase in the rate of S metabolism. Under the influence of I in the rate of metabolism of the inorganic form of S changed little, while that of the organic and not easily detachable form of S decreased more so at nitrate nutrition than at ammoniacal nutrition. By means of the paper chromatography method it was established that the amino acid composition of the root

Card 2/3

USSR/Soil Science. Mineral Fertilizers

J

Abs Jour : Ref Zhur-Biol., No 13, 1958, 58337, By N.N.
Sokolov

Author : Vlas'yuk P. A., and Butkevich A. P.
Inst : All-Union Academy of Agricultural Sciences imeni
V. I. Lenin
Title : Significance of Soil Microflora in the Manganese
Nutrition of Plants

Orig Pub : Dokl. VASKhNIL, 1957, No 5, 3-9

Abstract : The Ukrainian Scientific-Research Institute of
Plant Physiology conducted vegetation experiments
by growing oats, sugar beet, and flax with and
without rhizospheric microflora of these plants
in sandy cultures with the following variants:
a) without Mn, b) MnO_2 , c) $MnSO_4$ --1 norm; d).
[unclear]

Card 1/2

USSR/Soil Science. Mineral Fertilizers

J

Abs Jour : Ref Zhur-Biol., No 13, 1958, 58337, By N. N.
Sokolov

Abstract : $MnSO_4$ --10 norms. Rhizospheric microflora removed
the harmful effect of either the insufficiency
or excess of Mn and contributed to the mobility
of Mn in the variant with MnO_2 .

Editorial remark: $Fe(SO_4)_3$ and H_2BO_5 are mista-
kenly indicated as components of VNIS nutritive
medium. They should read $Fe_2(SO_4)_3$ and H_3BO_3 .

Card 2/2

USSR / Plant Physiology. Mineral Nutrition.

I-3

Abs Jour : Ref Zhur - Biol., No 10, 1958, No 43729

Author : Vlasuk, P. A.; Kosmatyy, Ye. S.; Klimovitskaya, Z. M.

Inst : Kiev Institute of Plant Physiology, AS USSR

Title : The Effect of Nitrate, Phosphorus, Potassium and Manganese Nutrients on Phosphorus Metabolism in the Sugar Beet.

Orig Pub : Izv. AN SSSR, Ser. biol., 1957, No. 5, 611-616

Abstract : A vegetative experiment (repeated five times) with the use of P^{32} , made at the Kiev Institute of Plant Physiology, showed that in the sugar beet culture Mn both on a nitrate ground and a ground of ammonium nitrogen nutrient increased the speed of the metabolism of P with RNA and DNA, as well as the P fraction of "nucleic acids plus phosphoproteins". The P metabolism speed of phospholipids and mineral phosphates was reduced under the influence of Mn on an ammonium nutrient ground and increased on a nitrate one. The P metabolism rate at a low phosphorus nutrient level reached

Card 1/2

VLASYUK, P.A.

USSR / Cultivated Plants. Plants for Technical Use. Oil
Plants. Sugar Plants.

M

Abs Jour : Ref Zhur - Biol., No 8, 1958, No 34771

Authors : Vlasuk, P. A.; Shmat'ko, I. G.
Inst : All-Union Scientific Research Institute for Sugar Beets
Title : Effects of Liquid Nitric Fertilizers on Seed Productivity.

Orig Pub : Sakharnaya svekla, 1957, No 6, 17-19

Abstract : Crop experiments conducted during the year 1956 by the All-Union Scientific Research Institute for Sugar Beets at the Kapitonovskiy sugar combine in the district of Cherkasskaya Oblast over an area of 18 hectares on medium-leached black earth ascertained the relative effects of liquid and solid nitric fertilizers, used on phosphate-potassic bases, on the productivity of sugar beet transplantation. To one hectare were added: potash salt (32% K₂O) 1 hwt; superphosphate 2.7 hwt; nitric acid of ammonia 1 hwt. All fer-

Card 1/2

USSR / Cultivated Plants. Plants for Technical Use. Oil M
Plants. Sugar Plants.

Abs Jour : Ref Zhur - Biol., No 18, 1958, No 34771

fertilizers were added as side-dressing. Liquid ammonia (82.3% N) was inserted into the soil by means of special machines to a depth of 12 cm. Liquid ammonia increased the soil acidity to a greater extent than nitric acid of ammonia. At the period of blooming, the intake of N into the balls and fruit stem leaves was most intensive under the action of the liquid ammonia; that of P, as a result of the added ammonium nitrate. The seed crop under the action of ammonia was by 1 to 2 hwgt per hectare larger than when using nitric acid ammonia. Thus, it could be concluded that liquid ammonia, when compared with all other nitric fertilizers, appears to be the best form of fertilization for the enriching of seed cultures of sugar beets. -- Smirnov.

Card 2/2

103

Country : USSR
Category : CULTIVATED PLANTS. POTATOES, Vegetables. Cucurbits.
Res. Jour. : ZEF ZHUR-SIOL., 21, 1953, NO. 65004
Author : Ulaev, P.A.; Derev'yanko, S.I.
Institut. : AS Ukrainian SSR
Title : The Effects of Different Forms of Potassium Fertilizers on the Physiologico-Biochemical Processes, Yield and Quality of Tomatoes and Potatoes Grown*
Orig. Pub. : Visnik AN URSR, 1957, No. 9, 42-52

Abstract : In tests conducted with the mid-season maturing Krasnodarats variety tomato and Lorkh potato under irrigation, an investigation has been made of the activity of ferment and respiration, the accumulation of ascorbic acid, sugar and chlorophyll, the change in moisture, as well as the Cl, S and K contents. K₂ and potassium-magnesium increased the vitamin C and dry matter content in tomato fruits from 4.8-5.0 (in the control) to

* Under Irrigation

Card: 1/3

Country : M
Category : CULTIVATED PLANTS. POTATOES,

Abs. Jour. : REF ZHUR-BIOL., 21, 1958, NO. 96004

Author :
Institut. :
Title :

Orig. Pub. :

Abstract : 5.4-5.8. All forms of K reduced the acidity of the fruit and favorably affected the water balance in the plants. The synthesis of chlorophyll in the tomato leaves and potato leaves increased only with the application of potash-magnesium and Kc. The latter produced the optimum respiration rate in the potato leaves. Potassium fertilization did not show any effect on a number of biochemical processes. Tomato yield boosts of 28.9, 26.8, and 18.2 cwt/ha. were obtained over the 226.2 of the

Card: 2/3

USSR / General Biology. Physical and Chemical
Biology.

B

Abs Jour : Ref Zhur - Biol., No 19, 1958, No. 35491

Author : Vlasyuk, P. A.

Inst : All-Union Academy of Agricultural Services imeni
V.I. Lenin.

Title : Basic Mechanisms of Biological Effects of Small
, Doses of Nuclear Irradiation.

Orig Pub : Dokl. VASKhNIL, 1957, No 10, 8-14

Abstract : A review of studies on the effects of irradia-
tion with radioactive isotopes on different a-
gricultural crops under conditions of vegetative
and field experiments, conducted by the Institu-
te of Plant Physiology of the Ukrainian Academy
of Agricultural Sciences, beginning in 1950.
With indicator doses of P^{32} , S^{35} , Ca^{45} , Zn^{65} and

Card 1/3

USSR / General Biology. Physical and Chemical
Biology.

B

Abs Jour : Ref Zhur - Biol., No 19, 1958, No 85491

Co^{60} , applied for pre-sowing treatment of seeds (by maceration in solutions of salts of these isotopes) or for extra root plant feeding at different stages, a considerable increase in productivity was attained in the great majority of the cases and in improvement in the quality of various crops (sugar beets, tomatoes, winter wheat, barley, rye, corn, clover, lupine, potatoes, grapes, tea, kok-saghyz). It was established in a number of experiments that even under unfavorable environmental conditions (lowered temperature, poor insulation) there is a positive effect of radioisotopes on plants. Investigations of the biological activity mechanism of ionizing radiations have shown that the responsive reaction of the living

Card 2/3

VLASYUK, P.A., akademik; DOBROTVORSKAYA, A.M., kandidat sel'skokhozynystvennykh nauk; GORDIYENKO, S.A.

Intensity of ferment action in the rhizosphere of various agricultural plants. Dokl. Akad. sel'skokhoz. 22 no.3:14-19 '57.

(MLRA 10:6)

1. Ukrainskiy nauchno-issledovatel'skiy institut fiziologii rasteniy.

(Rhizosphere microbiology) (Enzymes)

VLASYUK, P.A., akademik; BUTKEVICH, K.P.

Role of soil microflora in the manganese nutrition of plants.
Dokl.Akad.sel'khoz.22 no.5:3-9 '57. (MLRA 10:9)

1. Ukrainskiy nauchno-issledovatel'skiy institut fiziologii rasteniy.
(Rhizosphere microbiology) (Manganese) (Plants--Nutrition)

Vilasyuk, P.A.

VILASYUK, P.A., akademik.

Principle consistent manifestations of the biological action of
small doses of nuclear radiation. Dokl. Akad. sel'khoz. 22 no.10:
8-14 '57. (MIRA 10:12)

(Plants, Effect of radioactivity on)

VLASYUK, P.A.; DEREV'YANKO, S.I.

Effect of various potassium fertilizers on physiological and biochemical processes, yield and quality of tomatoes and potatoes in irrigation farming. Visnyk AN URSR 28 no.9:42-52 S '57.

(MIRA 11:1)

(Fertilizers and manures) (Irrigation farming)

Country : USSR I
Category : Plant Physiology. Mineral Nutrition.
Abs Jour. : Ref. Zbir.-Biologiya No. 11, 1956. No.48539
Author : Vlas'yuk, R.A.; Porutskiy, G.V.; Cherednichenko, S.V.
Institute : Acad. Sciences USSR
Title : Non-Root Side-Dressing with Thiamine and Plant
Growth during Florescence
Orig. Pub.: Dokl. AN SSSR, 1957, 112, No. 4, 769-771
Abstract : Top-dressing corn with thiamine during the flower-
ing period (in field experiments with both stable
and radioactive prepartes) stimulated the verti-
cal growth of the stalks and intensified the
growth of the reproductive organs during flower-
ing, yielding a slight boost in the grain harvest.
A direct correlation was noted between the plant
and the activity of the volatile organic secre-
tions of the plants, which was determined by
Card: 1/2

Country : USSR I
Category : Plant Physiology. Mineral Nutrition.

Abs. Jour.: Ref. Zhur.-Biolichya No. 11, 1958. No. 48539

Author :
Institute :
Title :

Orig. Pub.:

Abstract : measuring the gastic flexure they had produced.
--N.I. Borisova

Card: 2/2

Vlas'yuk, P. A.

USSR/Plant Physiology - General Problems.

I.

Abs Jour : Ref Zhur - Biol., No 18, 1958, 81961

Author : Vlas'yuk, P.A., Porutskiy, G.V., Cherednichenko, S.V.,
Dolgiy, S.N.

Inst : AS USSR

Title : The Influence of Extra-Root Fertilization on the Increase
of Germination of the Seed Material.

Orig Pub : Dokl. AN USSR, 1957, 113, No 1, 214-216

Abstract : Extra root fertilization with solutions of common (0,005%)
and radioactive (2.5 M curie per plant) thiamine of P^{31}
and P^{32} (10^{-4} M curie on a plant) was carried out by
spraying from an airplane or a tractor sprayer. The doses
were calculated on the basis of 200-400 l/ha. These ex-
periments took place in field and industrial tests dur-
ing the blossoming of corn and winter wheat.

Card 1/2

USSR/Plant Physiology - General Problems.

I.

Abs Jour : Ref Zhur - Biol., No 18, 1958, 81961

The dose of radioactive substances caused an increase in the energy of seed germination after the phase of wax ripeness. The reason for this increase is the earlier advent of physiological maturity as well as the shortening of the time of restoration of the changes in the bioelectrical potential (responding to a weak electrical stimulus), while conserving the capacity for synthesis of thiamine. These experiments were carried out at the Institute of Plant Physiology and Agricultural Chemistry at the AS UkrSSR. -- B.E. Kravtsova

Card 2/2

- 3 -

VLASYUK, Petr Antinovich [Vlesiuk, P.A.], akademik; SIROCHENKO, I.A.,
prof., red.; TUBOLEVA, M.V. [Tubolieva, M.V.], red.

[New microfertilizers] Novi mikrodobryva. Kyiv, 1958. 42 p.
(Tovarystvo dlia poshyrennia politychnykh i naukovykh znan'
Ukrains'koi RSR. Ser.3, no.8) (MIRA 12:3)
(Trace elements)

VLASYUK, P.A., akademik; ZEROV, D.K., akademik; PSHENICHNYY, P.D., akademik;
ROMANENKO, I.H., akademik, otvetstvennyy red.; MOVCHAN, V.A.;
RODIONOV, S.P.; TYULENEV, N.A.; DAVYDOV, G.M., kand. ekon. nauk;
KUGUKALO, I.A., kand. ekon. nauk; BEREZIKOV, V.S.; FEDUN, A.D.;
GRUDZINSKAYA, O.S., red. izd-va; YURCHISHIN, V.I., tekhn. red.

[Natural conditions and resources of the Polesye; transactions of
the Conference on Problems of the Development of the Productive
Forces of the Ukrainian Polesye] Prirodnye usloviia i resursy
Poles'ia; trudy konferentsii po voprosam razvitiia proizvoditel'-
nykh sil Poles'ia USSR. Kiev. Pt.1. 1958. 123 p. (MIRA 11:?)

1. Akademiya nauk URSR, Kiev. Rada po vyzchenniu produktivnykh syl.
2. Akademiya nauk USSR (for Vlasyuk, Zerov). 3. Ukrainskaya
akademiya sel'skokhozyaystvennykh nauk (for Vlasyuk, Pshenichnyy,
Romanenko). 4. Vsesoyuznaya akademiya sel'skokhozyaystvennykh nauk
imeni V.I. Lenina (for Vlasyuk). 5. Chlen-korrespondent Vsesoyuz-
noy akademii sel'skokhozyaystvennykh nauk imeni V.I. Lenina (for
Romanenko). 6. Chlen-korrespondent akademii nauk USSR (for Movchan,
Rodionov, Tyulenev). 7. Zamestitel' nachal'nika otdela svodnykh
perspektivnykh planov Gosplana USSR (for Berezikov). 8. Nachal'nik
podotdela sel'skogo khozyaystva otdela svodnykh perspektivnykh planov
Gosplana USSR (Fedun).

(Polesye--Natural resources)

ROMANENKO, I.N., akademik, otvetstvennyy red.; VLASYUK, P.A., akademik, red.; ZEROV, D.K., akademik, red.; RODIONOV, S.P., red.; TYULENEV, N.A., red.; PSHENICHNYY, P.D., akademik, red.; DAVYDOV, G.M., kand. ekon. nauk, red.; KUGUKALO, I.A., kand. ekon. nauk, red.; BEREZIKOV, V.S., red.; FEDUN, A.D., red.; KOZAKEVICH, T.A., red. izd-va; SIVACHENKO, Ye. K., tekhn. red.

[Problems in the economy of Polesye; transactions of a conference]
Voprosy ekonomiki Poles'ia; trudy konferentsii. Kiev, Izd-vo Akad. nauk USSR. Vol. 4. 1958. 134 p. (MIRA 11:10)

1. Konferentsiya po voprosam razvitiya proizvoditel'nykh sil Poles'ya USSR. 1955.
2. Akademiya nauk USSR (for Vlasyuk, Zerov.).
3. Ukrainskaya Akademiya sel'skokhozyaystvennykh nauk (for Vlasyuk, Romanenko, Pshenichnyy).
4. Vsesoyuznaya Akademiya sel'skokhozyaystvennykh nauk im. V.I.Lenina (for Vlasyuk).
5. Chlen-korrespondent Vsesoyuznoy Akademii sel'skokhozyaystvennykh nauk im. V.I.Lenina (for Romanenko).
6. Chlen-korrespondent Akademii nauk USSR (for Rodionov, Tyulenev).
7. Zamestitel' nachal'nika ot dela svodnykh perspektivnykh planov Gosplana Soveta Ministrov USSR (for Berezikov).
8. Nachal'nik podot dela sel'skogo khozyaystva i zagotovok ot dela svodnykh perspektivnykh planov sel'skogo khozyaystva Gosplana Soveta Ministrov USSR (for Fedun).

(Polesye--Economic conditions)

VLASYUK, P.A., akademik, red.; ROMANENKO, I.N., akademik, red.; RODIONOV, S.P., red.; TYULENEV, red.; PSHENICHNYY, P.D., akademik, red.; DAVYDOV, kand.ekon.nauk, red.; KUGUKALO, I.A., kand.ekon.nauk; BEREZIKOV, V.S., red.; FEDIN, A.D., red.; KOZAKEVICH, T.A., red. izd-va; SIVACHENKO, Ye.K., tekhn.red.

[Proceedings of the Conference on Problems in Developing Production in Polesye] Konferentsiya po voprosam razvitiia proizvoditel'nykh sil Poles'ia USSR. Kiev, 1955. Pt.3 [Problems in the development of agriculture in Polesye; stockbreeding and feed supply, land improvement and reclamation of swamps] Voprosy razvitiia sel'skogo khoziaistva Poles'ia; zhivotnovodstvo i kormovaia baza, melioratsiya i osvoenie bolot. Kiev, Izd-vo Akad. nauk USSR. 1958. 208 p. (MIRA 12:1)

1. AN USSR; Ukrainskaya akademiya sel'skokhoz.nauk i Vsesoyuznaya akademiya sel'skokhoz.nauk im. V.I. Lenina (for Vlasyuk). 2. Ukrainskaya akademiya sel'skokhoz.nauk, chlen-korrespondent Vsesoyuznoy akademii sel'skokhoz. nauk im. V.I. Lenina (for Romanenko). 3. Chlen-korrespondent AN USSR (for Rodionov, Tyulenev). 4. Institut fizioligii rasteniy i agrokhimii AN USSR (for Tyulenev). 5. Ukrainskaya akademiya sel'skokh. nauk (for Psheinichnyy). 6. Zamestitel' nachal'nika ot dela svodnykh perspektivnykh planov Gosplana USSR (for Berezikov); 7. Nachal'nik podot dela sel'skogo khozyaystva ot dela svodnykh perspektivnykh planov Gosplana USSR (for Fedin).

(Polesye--Agriculture)

VLASYUK, PAV.

3/7

PHASE I BOOK EXPLOITATION

SOV/233

Konferentsiya po agrometeorologii i agroklimatologii Ukrainskoj SSR
 Materialy konferentsii. (Material of the Conference on Agricultural Meteorology and Climatology of the Ukrainian SSR). Leningrad.
 Glidrometeorologicheskij izdatel'stvo. 1958. 247 p. Errata slip inserted.
 printed.

Sponsoring Agencies: USSR. Glavnaya upravleniya Gidrometeorologicheskoy sluzhby Ukrainskoj SSR. Ministerstvo sel'skogo khozyaistva, Ukrainskiy nauchno-issledovatel'skiy gidrometeorologicheskiy in-tstitut, i Ukrainskaya akademiya sel'skokhozyaistvennykh nauk.
 Resp. Ed.: O.P. Pril'hot'ko; Ed.: V.D. Pisarevskaya; Tech. Ed.: K.I. Braymina.

PURPOSE: This book is intended for agriculturists, agrometeorologists, and instructors in related areas.

COVERAGE: This collection of articles deals with problems in agricultural meteorology in the Ukraine. Among the topics discussed are: wintering, planting time for winter crops, corn cultivation, potato degeneration, moisture supply, and adverse weather factors.

References accompany individual articles.

TABLE OF CONTENTS:

Baev, I.I. [Deputy Minister of Agriculture, Ukr. SSR]	Introductory Word	3
Bozatyry, T.K. [Chief of the Hydroagrometeorological Service, Ukr. SSR]	Practical Hydroagrometeorological Service for Agricultural Production in the Ukraine	5
Kolouch, A.M. [Ukrainian Scientific Research Hydromet-Institute]	Regional Agroclimatological (Reference Books) of the Ukraine and Their Application in Production	10
Pril'hot'ko, O.P. [Ukrainian Scientific Research Hydromet-Institute]	The State of Agrometeorological Studies in the Ukraine	15
Konashchukova, M.N. [Ukrainian Scientific Research Hydromet-Institute]	Organization and Utilization of Meteorological Observations of Departmental Stations in Scientific Work and Agricultural Practice	20
Vlas'yuk, P.A. and M.A. Gurileva [Ukrainian Scientific Research Institute for Plant Physiology]	Special Features of the Wintertime Over of Winter Crops in 1955-56 in Various Regions of the USSR	23
Lishikaki, V.M. [Ukrainian Scientific Research Hydromet-Institute]	Agrometeorological Conditions of the Wintering of Winter Crops in the Ukraine	31
Lishikaki, V.M. [Ukrainian Scientific Research Hydromet-Institute]	Agroclimatic Basis for the Planting Time of Winter Crops in the UkrSSR	36
Mil'ner, Ye. S. [Central Institute of Prognoses]	Relationship Between the Phases in the Development of Winter Crops in Autumn and the Agrometeorological Conditions. Probability of Climate Development of Winter Crops as Related to the Different Planting Time in the Ukraine	69

SOV/2384

PIASHE I BOOK EXPLOITATION

317)

Konferentsiya po agroeteorologii i agroklimatologii: Ukrainskoy SSR
Materialy konferentsii (Material of the Conference on Agricultural
Meteorology and Climatology of the Ukrainian SSR) Leningrad,
Ukrainskii nauchno-issledovatel'skiy gidrometeorologicheskiy in-
stitut, and Ukrainskaya akademiya zemel'noy i gospodarki sel'skogo
prirodozrzadstva, 1958. 257 p. Errone slip inserted. 700 copies
printed.

Sponsoring Agencies: USSR. Glavnoye upravleniye gidrometeorologicheskoy sluzhby. Ukrainian SSR. Ministerstvo sel'skogo khozyaystva.
Ukrainskii nauchno-issledovatel'skiy gidrometeorologicheskiy in-
stitut, and Ukrainskaya akademiya sel'skokhozyaystvennykh nauk.
Resp. Ed.: G.P. Fridhot'ko; Ed.: V.D. Pisarevskaya; Tech. Ed.:
M.I. Bravina.

PURPOSE: This book is intended for agriculturists, agrometeorolo-
gists, and instructors in related areas.

COVERAGE: This collection of articles deals with problems in agricultural
cultural meteorology in the Ukraine. Among the topics discussed
are: winterizing, planting time for winter crops, corn cultivation,
potato degeneration, moisture supply, and adverse weather factors.
References accompany individual articles.

TABLE OF CONTENTS:

- Fedorova, N.A. [Ukrainian Scientific Research Institute for Agri-
culture] Significance of Planting Time for the Wintering of Winter
Crops Under Poles'e (Woodlands) and Northern Lesotep (Forested
Steppe Regions) Conditions in the USSR 76
Michurinskaya, M.I. [Ukrainian Scientific Research Institute of Crop
Science] Significance of Critical Temperatures in Forecasting the
Wintering Conditions 84
Ovrlivka, M.A. [Ukrainian Scientific Research Institute for Plant
Physiology] Forecasting the Reaction of the Various Grades of Winter
Wheat Upon the Intermittent Temperatures of the Winter and Early
Spring Periods 91
Ovrlivka, M.A. and N.A. Fedorova. Results of Checking the Method
for Determining the Viability of Winter Crops by the Conditions of
the Vegetative Cone 96
Iovenko, M.G. [Ukrainian Scientific Research Hydromet Institute]
Moisture Reserves of Various Climatic Soil Zones of the Ukraine 100
Zemets, O.M. [All-Union Scientific Research Institute for Study of
card 4/7

1/2

USM/Soil Science. Organic Fertilizers

J-6

Ree Jour : Ref Zhur - Biol., № 20, 1958, № 91486

Author : Vlas'yuk P.I., Darmenko M.S., Monorik A.V.

Inst

Title : The Effectiveness of the Use of Industrial Brown Coal Waste
for Various Agricultural Crops

Orig Pub : Byul. po fiziol. rasteniy, 1958, No 2, 48-52

Abstract : No abstract

E N D

Card : J/1

39

VLASYUK, P.A.; LENDENSKAYA, L.D.

Polar distribution of manganese in different parts of organs of wheat and corn plants [with summary in English]. Fiziol.rast. 5 no.6: 488-493 N-D '58. (MIRA 11:12)

1. Institut fiziologii rasteniy Ukrainskoy SSR, Kiiev.
(Polarity (Biology)) (Plants, Effect of manganese on)
(Plants, Motion of fluids in)

VLASYUK, P.A., akademik; GURIL'OVA, M.A. [Guryl'ova, M.A.], kand.biol.
nauk

Winter hardiness of plants. Nauka i zhyttia 8 no.3:23-25
Mr '58. (MIRA 12:9)

1. AN USSR i Vsesoyuznaya akademiya sel'skokhozyaystvennykh nauk
im. Lenina (for Vlasyuk).
(Plants--Frost resistance)

SOV-21-58-8-22/27

AUTHORS: Vlasyuk, P.A., Member of the AS UkrSSR, and Lisoval, P.Z.

TITLE: Effect of Various Forms of Potassium Fertilizers on the Yield of Farm Crops (Vliyanije razlichnykh form kaliynykh udobreniy na urozhay sel'skokhozyaystvennykh kul'tur)

PERIODICAL: Dopovidi Akademii nauk Ukrains'koj RSR, 1958, Nr 8,
pp 887-890 (USSR)

ABSTRACT: With the aim of establishing the best conditions of nutrition and in raising the yield of plants, the authors studied various forms of potassium fertilizers from deposits of the western regions of the Ukrainian SSR on permanent fields of the Ukrainian Research Institute for Plant Physiology and temporary fields at the Irpen' river and at the Kherson research station. As a result of many years of research, they arrived at the conclusion that to obtain high yields and a better quality of farm produce, both for grain and technical as well as for vegetable and fodder crops, it is essential to utilize not only chloride but sulfate and sulfate-magnesian forms of potassium fertilizers and their combinations.

There are 6 Soviet references.

Card 1/2